


Cat. No:	MAB-94617
Size:	100 ug
Clone:	PR10
Concentration:	1mg/ml
Host:	Ms
Isotype:	IgG
Immunogen:	Recombinant protein of human PRKACA.
Reactivity:	Hu, Ms, Rt
Applications:	Western Blot: 1:500 - 1:2000 Immunofluorescence: 1:50 - 1:200
Molecular Weight:	40kDa
Purification:	Aff. Pur.
Synonyms:	PRKACA; PKACA; PPNAD4; cAMP-dependent protein kinase catalytic subunit alpha


This gene encodes one of the catalytic subunits of protein kinase A, which exists as a tetrameric holoenzyme with two regulatory subunits and two catalytic subunits, in its inactive form. cAMP causes the dissociation of the inactive holoenzyme into a dimer of regulatory subunits bound to four cAMP and two free monomeric catalytic subunits. Four different regulatory subunits and three catalytic subunits have been identified in humans. cAMP-dependent phosphorylation of proteins by protein kinase A is important to many cellular processes, including differentiation, proliferation, and apoptosis. Constitutive activation of this gene caused either by somatic mutations, or genomic duplications of regions that include this gene, have been associated with hyperplasias and adenomas of the adrenal cortex and are linked to corticotropin-independent Cushing's syndrome. Alternative splicing results in multiple transcript variants encoding different isoforms. Tissue-specific isoforms that differ at the N-terminus have been described, and these isoforms may differ in the post-translational modifications that occur at the N-terminus of some isoforms.

Background:

Form:	Liquid
Buffer:	PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Storage:	Store at -20°C. Avoid freeze / thaw cycles.



Western blot analysis of extracts of various cell lines, using PRKACA antibody at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-



Immunofluorescence analysis of BALB-3T3 cells using PRKACA Mouse Monoclonal Antibody

Mouse IgG (H+L) at 1:10000 dilution.
Lysates/proteins: 25ug per lane.
Blocking buffer: 3% nonfat dry milk in
TBST.
Detection: ECL Basic Kit.
Exposure time: 1s.

at dilution of 1:100 (40x lens).
Blue: DAPI for nuclear staining.