

Cat. No:	ABN16535
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human PCDH-X/Y. AA range:531-580
Reactivity:	Human,Rat,Mouse
Applications:	IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000
Purification:	Affinity purification
Synonyms:	PCDH11Y; PCDH11; PCDH22; PCDHY; Protocadherin-11 Y-linked; Protocadherin-11; Protocadherin on the Y chromosome; PCDH-Y; Protocadherin prostate cancer; Protocadherin-PC; Protocadherin-22; PCDH11X; KIAA1326; PCDH11; PCDHX; Protocadherin-11 X-
Background:	<p>This gene belongs to the protocadherin family, a subfamily of the cadherin superfamily. The encoded protein consists of an extracellular domain containing seven cadherin repeats, a transmembrane domain, and a cytoplasmic tail that differs from those of the classical cadherins. This gene is located on the Y chromosome in a block of X/Y homology and is very closely related to its paralog on the X chromosome. The protein is thought to play a role in cell-cell recognition during development of the central nervous system. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2013],alternative products:Additional isoforms seem to exist,disease:A chromosomal aberration involving PCDH11Y is a cause of multiple congenital abnormalities, including severe bilateral vesicoureteral reflux (VUR) with ureterovesical junction defects. Translocation t(Y;3)(p11;p12) with ROBO2.,function:Potential calcium-dependent cell-adhesion protein.,similarity:Contains 7 cadherin domains.,subunit:Interacts with CTNNB1.,tissue specificity:Expressed strongly in fetal brain and brain (cortex, amygdala, thalamus, substantia nigra, hippocampus, caudate nucleus and corpus callosum). Expressed at low level in testis. Expressed in apoptosis-resistant cells.,</p>
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only
IMMUNOLOGICAL SCIENCES**