

Cat. No:	ABN16545
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human PRPF18. AA range:182-231
Reactivity:	Human,Mouse,Rat
Applications:	WB 1:500-1:2000,IHC 1:50-1:300,ELISA 1:2000-1:20000
Molecular Weight:	40kDa
Purification:	Affinity purification
Synonyms:	PRPF18; HPRP18; Pre-mRNA-splicing factor 18; PRP18 homolog; hPRP18
Background:	Pre-mRNA splicing occurs in 2 sequential transesterification steps. The protein encoded by this gene is found to be essential for the catalytic step II in pre-mRNA splicing process. It is found in the spliceosome, and contains seven WD repeats, which function in protein-protein interactions. This protein has a sequence similarity to the yeast splicing factor Prp18. [provided by RefSeq, Jul 2008],function:Participates in the second step of pre-mRNA splicing.,similarity:Belongs to the PRP18 family.,subcellular location:Colocalizes with spliceosomal snRNPs.,subunit:Heterodimer with PPIH. Interacts with PRPF4 and with the spliceosome. Part of a complex containing U4/U6 snRNPs.,
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only
IMMUNOLOGICAL SCIENCES**