

|                          |  |
|--------------------------|--|
| <b>Cat. No:</b>          | AB-84762   |
| <b>Conjugate:</b>        | Unconjugated   |
| <b>Size:</b>             | 100 ug   |
| <b>Clone:</b>            | POLY   |
| <b>Concentration:</b>    | 1mg/ml   |
| <b>Host:</b>             | Rabbit   |
| <b>Isotype:</b>          | IgG  |
| <b>Immunogen:</b>        | The antiserum was produced against synthesized peptide derived from human SHP-1. AA range:502-551  |
| <b>Reactivity:</b>       | Human, Mouse, Rat  |
| <b>Applications:</b>     | Western Blot: 1/500 - 1/2000 Immunohistochemistry: 1/100 - 1/300<br>Immunofluorescence: 1:50-200 ELISA: 1/20000  |
| <b>Molecular Weight:</b> | 67kD   |
| <b>Purification:</b>     | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography on   |
| <b>Synonyms:</b>         | PTPN6; HCP; PTP1C; Tyrosine-protein phosphatase non-receptor type 6;<br>Hematopoietic cell protein-tyrosine phosphatase; Protein-tyrosine phosphatase<br>1C; PTP-1C; Protein-tyrosine phosphatase SHP-1; SH-PTP1   |
| <b>Background:</b>       | The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. N-terminal part of this PTP contains two tandem Src homolog (SH2) domains, which act as protein phospho-tyrosine binding domains, and mediate the interaction of this PTP with its substrates. This PTP is expressed primarily in hematopoietic cells, and functions as an important regulator of multiple signaling pathways in hematopoietic cells. This PTP has been shown to interact with, and dephosphorylate a wide spectrum of phospho-proteins involved in hematopoietic cell signaling. Multiple alternatively spliced variants of this gene, which encode distinct isoforms, have been reported. |
| <b>Form:</b>             | Liquid   |
| <b>Buffer:</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| <b>Storage:</b>          | Store at -20°C. Avoid repeated freeze-thaw cycles.   |

**For Research use only  
IMMUNOLOGICAL SCIENCES**