

Cat. No:	AB-84321
Size:	100ug
Clone:	POLY
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgG
Immunogen:	A synthetic peptide corresponding to a sequence of human Tbet/ Tbx21(NNVTQMIVLQSLHKYQPRRHIVEVNDGE).
Reactivity:	Hu, Ms, Rt
Applications:	Western blot: 0.25-0.5ug/ml Flow Cytometry: 1-3ug/1×10 ⁶ cells,
Molecular Weight:	58kDa
Purification:	Aff. Pur.

Background: T-box transcription factor TBX21 is a protein that in humans is encoded by the TBX21 gene. It is mapped to 17q21.32. This gene is a member of a phylogenetically conserved family of genes that share a common DNA-binding domain, the T-box. T-box genes encode transcription factors involved in the regulation of developmental processes. This gene is the human ortholog of mouse Tbx21/Tbet gene. Studies in mouse show that Tbx21 protein is a Th1 cell-specific transcription factor that controls the expression of the hallmark Th1 cytokine, interferon-gamma (IFNG). Expression of the human ortholog also correlates with IFNG expression in Th1 and natural killer cells, suggesting a role for this gene in initiating Th1 lineage development from naive Th precursor cells.

Form: Liquid

Buffer: Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃.

Storage: At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.





Figure 1. Western blot analysis of TBX21 using anti-TBX21 Antibody.
Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded



Immunofluorescent analysis of 293 cells labeling T-bet / Tbx21 with AB-84247 at 5 µg/ml.




Figure 2. Flow Cytometry analysis of HEPA1-6 cells using anti-TBX21 antibody. Overlay histogram showing HEPA1-6 cells stained with (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-TBX21 Antibody 1µg/1×10⁶ cells) for 30 min at

with 50ug of sample under reducing conditions.
Lane 1: mouse RAW246.7 whole cell lysates.
After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes.
Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TBX21 antigen affinity purified polyclonal antibody at 0.5 mug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT.
The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system.
A specific band was detected for TBX21 at approximately 65KD. The expected band size for TBX21 is at 58KD.

20°C. DyLight®488 conjugated goat anti-rabbit IgG (5-10mug/1×10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1mug/1×10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




Figure 3. Flow Cytometry analysis of NRK cells using anti- TBX21 antibody. Overlay histogram showing NRK cells stained with (Blue line).The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-TBX21 Antibody (1mug/1×10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (5-10mug/1×10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1mug/1×10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control




Figure 4. Flow Cytometry analysis of mouse PBMC cells using anti-TBX21 antibody Overlay histogram showing mouse PBMC cells stained with (Blue line).The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-TBX21 Antibody (1mug/1×10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (5-10mug/1×10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1mug/1×10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.