

|                       |   |
|-----------------------|---|
| <b>Cat. No:</b>       | ABN18893  |
| <b>Conjugate:</b>     | Unconjugated  |
| <b>Size:</b>          | 100µL   |
| <b>Clone:</b>         | Polyclonal  |
| <b>Concentration:</b> | 1mg/ml  |
| <b>Host:</b>          | Rabbit  |
| <b>Isotype:</b>       | IgG   |
| <b>Immunogen:</b>     | The antiserum was produced against synthesized peptide derived from human THBS4. AA range:561-610   |
| <b>Reactivity:</b>    | Human,Rat,Mouse   |
| <b>Applications:</b>  | WB 1:500-1:2000,IHC 1:100-1:500,ICC/IF 1:100-1:500,ELISA 1:5000-1:20000   |
| <b>Purification:</b>  | Affinity purification   |
| <b>Synonyms:</b>      | THBS4; TSP4; Thrombospondin-4   |
| <b>Background:</b>    | <p>thrombospondin 4(THBS4) Homo sapiens The protein encoded by this gene belongs to the thrombospondin protein family. Thrombospondin family members are adhesive glycoproteins that mediate cell-to-cell and cell-to-matrix interactions. This protein forms a pentamer and can bind to heparin and calcium. It is involved in local signaling in the developing and adult nervous system, and it contributes to spinal sensitization and neuropathic pain states. This gene is activated during the stromal response to invasive breast cancer. It may also play a role in inflammatory responses in Alzheimer's disease. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Apr 2015],function:Adhesive glycoprotein that mediates cell-to-cell and cell-to-matrix interactions. Can bind to fibrinogen, fibronectin, laminin and type V collagen.,similarity:Belongs to the thrombospondin family.,similarity:Contains 1 TSP C-terminal (TSPC) domain.,similarity:Contains 1 TSP N-terminal (TSPN) domain.,similarity:Contains 4 EGF-like domains.,similarity:Contains 8 TSP type-3 repeats.,subunit:Homopentamer; disulfide-linked.,</p> |
| <b>Form:</b>          | Liquid  |
| <b>Buffer:</b>        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.  |
| <b>Storage:</b>       | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.  |

**For Research use only  
IMMUNOLOGICAL SCIENCES**