

<b>Cat. No:</b>	MABN83717
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Monoclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	A synthesized peptide derived from human TATA element modulatory factor
<b>Reactivity:</b>	Human
<b>Applications:</b>	WB 1:1000-1:2000, ICC 1:50-1:200
<b>Molecular Weight:</b>	Calculated MW: 123 kDa ; Observed MW: 160 kDa
<b>Purification:</b>	Affinity Purification
<b>Synonyms:</b>	ARA160; TMF; Tmf1;;TATA element modulatory factor
<b>Background:</b>	Potential coactivator of the androgen receptor. Mediates STAT3 degradation. May play critical roles in two RAB6-dependent retrograde transport processes: one from endosomes to the Golgi and the other from the Golgi to the ER. This protein binds the HIV-1 TATA element and inhibits transcriptional activation by the TATA-binding protein (TBP).
<b>Form:</b>	Liquid
<b>Buffer:</b>	Purified antibody in PBS with 0.05% sodium azide, 0.05% BSA and 50% glycerol.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only  
IMMUNOLOGICAL SCIENCES**