

Cat. No:	ABN19072
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human TMPRSS3. AA range:405-454
Reactivity:	Human,Mouse
Applications:	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000
Molecular Weight:	49kDa
Purification:	Affinity purification
Synonyms:	TMPRSS3; ECHOS1; TADG12; Transmembrane protease serine 3; Serine protease TADG-12; Tumor-associated differentially-expressed gene 12 protein
Background:	<p>This gene encodes a protein that belongs to the serine protease family. The encoded protein contains a serine protease domain, a transmembrane domain, an LDL receptor-like domain, and a scavenger receptor cysteine-rich domain. Serine proteases are known to be involved in a variety of biological processes, whose malfunction often leads to human diseases and disorders. This gene was identified by its association with both congenital and childhood onset autosomal recessive deafness. This gene is expressed in fetal cochlea and many other tissues, and is thought to be involved in the development and maintenance of the inner ear or the contents of the perilymph and endolymph. This gene was also identified as a tumor-associated gene that is overexpressed in ovarian tumors. Alternatively spliced transcript variants have been described. [provided by RefSeq, Jan 2012],disease:Defects in TMPRSS3 are a cause of non-syndromic sensorineural deafness autosomal recessive type 10 (DFNB10) [MIM:605316].,disease:Defects in TMPRSS3 are the cause of non-syndromic sensorineural deafness autosomal recessive type 8 (DFNB8) [MIM:601072]. DFNA8 is a form of sensorineural hearing loss. Sensorineural deafness results from damage to the neural receptors of the inner ear, the nerve pathways to the brain, or the area of the brain that receives sound information.,function:Probable protease. Seems to be capable of activating ENaC.,PTM:Undergoes autoproteolytic activation.,similarity:Belongs to the peptidase S1 family.,similarity:Contains 1 LDL-receptor class A domain.,similarity:Contains 1 peptidase S1 domain.,similarity:Contains 1 SRCR domain.,tissue specificity:Expressed in many tissues including fetal cochlea. Isoform T is found at increased levels in some carcinomas.,</p>
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only
IMMUNOLOGICAL SCIENCES**