

<b>Cat. No:</b>	MABN02717
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Monoclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	A synthetic peptide of human TRIM23
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Applications:</b>	WB 1:500-1:1000
<b>Molecular Weight:</b>	Calculated MW: 64 kDa; Observed MW: 64 kDa
<b>Purification:</b>	Affinity Purification
<b>Synonyms:</b>	ARD1; ARFD1; RNF46
<b>Background:</b>	Acts as an E3 ubiquitin-protein ligase. Plays an essential role in autophagy activation during viral infection. Mechanistically, activates TANK-binding kinase 1/TBK1 by facilitating its dimerization and ability to phosphorylate the selective autophagy receptor SQSTM1. In order to achieve this function, TRIM23 mediates 'Lys-27'-linked auto-ubiquitination of its ADP-ribosylation factor (ARF) domain to induce its GTPase activity and its recruitment to autophagosomes (PubMed:28871090).
<b>Form:</b>	Liquid
<b>Buffer:</b>	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only  
IMMUNOLOGICAL SCIENCES**