

Cat. No:	MABN84310
Conjugate:	Unconjugated
Size:	100µL
Clone:	Monoclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	A synthesized peptide derived from human TRIM29
Reactivity:	Human, Mouse, Rat
Applications:	WB 1:1000-1:2000
Molecular Weight:	Calculated MW: 66 kDa ; Observed MW: 68 kDa
Purification:	Affinity Purification
Synonyms:	ATDC; TRIM29;;TRIM29
Background:	Plays a crucial role in the regulation of macrophage activation in response to viral or bacterial infections within the respiratory tract. Mechanistically, TRIM29 interacts with IKBKG/NEMO in the lysosome where it induces its 'Lys-48' ubiquitination and subsequent degradation. In turn, the expression of type I interferons and the production of pro-inflammatory cytokines are inhibited. Additionally, induces the 'Lys-48' ubiquitination of STING1 in a similar way, leading to its degradation.
Form:	Liquid
Buffer:	Purified antibody in PBS with 0.05% sodium azide, 0.05% BSA and 50% glycerol.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only
IMMUNOLOGICAL SCIENCES**