




Cat. No:	ABS152667
Conjugate:	unconjugated
Size:	200 ug
Clone:	POLY
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG, κ
Immunogen:	Recombinant protein corresponding to Human beta Tubulin
Reactivity:	Species, Human, Mouse, Rat
Applications:	Applications Dilution WB 1: 5000-1: 10000 ICC/IF 1: 100-1: 300
Purification:	Affinity purification
Synonyms:	Tubb5, Beta 5 tubulin, Class I beta tubulin, Tubulin beta class I, TUBB1, TUBB, TUBB5, Tubulin Beta-5 Chain, β tubulin, CDCBM6
Background:	Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha chain.
Form:	Liquid
Buffer:	PBS with 0.15% ProClin300, 100 μ g/mL BSA and 50% glycerol.
Storage:	Store at -20°C for one year. Avoid repeated freeze/thaw cycles.



WB analysis of beta Tubulin. Sample: Protein C6, were ICC analysis of beta Tubulin. Sample: C6, were ICC analysis of beta Tubulin. Sample: C6, were treated by RIPA Lysis Buffer. Blocking buffer: 3% fixed in pre-cooled methanol 10 minutes. Blocking buffer: 3% BSA in PBS, RT, 30min. Primary antibody: Nonfat dry milk in TBST, RT, 1h. Primary antibody: buffer: 3% BSA in PBS, RT, 30min. Primary antibody: 1: 5000, 4°C overnight. Secondary antibody: HRP antibody: 1: 100, 4°C overnight. Secondary antibody: 1: 300, 4°C overnight. Secondary conjugated Goat Anti-Rabbit IgG, 1: 5000, RT, 1h. antibody: Cy3 conjugated Goat Anti-Rabbit IgG, 1: 300, RT, 1h. antibody: Cy3 conjugated Goat Anti-Rabbit IgG, 1: 300, RT, 1h.

**For Research use only
IMMUNOLOGICAL SCIENCES**