

Cat. No:	MABN19471
Conjugate:	Unconjugated
Size:	100µL
Clone:	Monoclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	A synthetic peptide corresponding to target protein
Reactivity:	Human,Mouse,Rat
Applications:	WB 1:500-1:2000,IHC 1:50-1:200,ICC/IF 1:100-1:200,FC 1:20-1:50
Molecular Weight:	59kDa
Purification:	Affinity purification
Synonyms:	EC 1.14.16.2; TH isoform 3; TH isoform a, TH-4;TY3H; TYH; Tyrosine 3-hydroxylase; Tyrosine 3-monoxygenase; tyrosine hydroxylase;
Background:	Tyrosine hydroxylase (EC 1.14.16.2) is involved in the conversion of phenylalanine to dopamine. As the rate-limiting enzyme in the synthesis of catecholamines, tyrosine hydroxylase has a key role in the physiology of adrenergic neurons. Plays an important role in the physiology of adrenergic neurons (By similarity). Positively regulates the regression of retinal hyaloid vessels during postnatal development (By similarity).
Form:	Liquid
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only
IMMUNOLOGICAL SCIENCES**