

|                          |  |
|--------------------------|--|
| <b>Cat. No:</b>          | MABN84224  |
| <b>Conjugate:</b>        | Unconjugated   |
| <b>Size:</b>             | 100µL  |
| <b>Clone:</b>            | Monoclonal   |
| <b>Concentration:</b>    | 1mg/ml   |
| <b>Host:</b>             | Rabbit   |
| <b>Isotype:</b>          | IgG  |
| <b>Immunogen:</b>        | A synthesized peptide derived from human USP9X   |
| <b>Reactivity:</b>       | Human,Mouse,Rat  |
| <b>Applications:</b>     | WB 1:1000-1:2000,IHC 1:100-1:200,ICC/IF 1:50-1:200,ICC 1:50-1:200,FC 1:20-1:100  |
| <b>Molecular Weight:</b> | Calculated MW: 290 kDa ; Observed MW: 292 kDa  |
| <b>Purification:</b>     | Affinity Purification  |
| <b>Synonyms:</b>         | DDFRX; FAF; FafI; Fam; hFAM; MRX99; Usp9x; X chromosome; X-linked;;USP9X   |
| <b>Background:</b>       | Deubiquitinase involved both in the processing of ubiquitin precursors and of ubiquitinated proteins.May therefore play an important regulatory role at the level of protein turnover by preventing degradation of proteins through the removal of conjugated ubiquitin. |
| <b>Form:</b>             | Liquid   |
| <b>Buffer:</b>           | Purified antibody in PBS with 0.05% sodium azide,0.05% BSA and 50% glycerol.   |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.   |

**For Research use only  
IMMUNOLOGICAL SCIENCES**