




Cat. No:	AB-82183
Conjugate:	Unconjugated
Size:	200 ug
Clone:	POLY
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human VEGF-A. AA range:110-159
Reactivity:	Human, Mouse, Rat, Pig, Rabbit
Applications:	Western Blot: 1/500 - 1/2000 Immunofluorescence: 1:50-200 Immunohistochemistry: 1/100 - 1/300 ELISA:1/10000.
Molecular Weight:	21kD(monomer),42kD(dimer)
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Synonyms:	VEGFA; VEGF; Vascular endothelial growth factor A; VEGF-A; Vascular permeability factor; VPF
Background:	This gene is a member of the PDGF/VEGF growth factor family. It encodes a heparin-binding protein, which exists as a disulfide-linked homodimer. This growth factor induces proliferation and migration of vascular endothelial cells, and is essential for both physiological and pathological angiogenesis. Disruption of this gene in mice resulted in abnormal embryonic blood vessel formation. This gene is upregulated in many known tumors and its expression is correlated with tumor stage and progression. Elevated levels of this protein are found in patients with POEMS syndrome, also known as Crow-Fukase syndrome. Allelic variants of this gene have been associated with microvascular complications of diabetes 1 (MVCD1) and atherosclerosis. Alternatively spliced transcript variants encoding different isoforms have been described.
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. and 0.02% sodium azide.
Storage:	Store at -20°C. Avoid repeated freeze-thaw cycles.



Immunohistochemical analysis of paraffin-embedded Human skeletal



Immunofluorescence analysis of A549.

Western blot analysis of various lysates,
using VEGFA Rabbit pAb at 1:1000
dilution.

Secondary antibody: HRP Goat Anti-
Rabbit IgG (H+L) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.

Blocking buffer: 3% nonfat dry milk in
TBST.

Detection: ECL West Pico Plus.

Exposure time: 30s.

muscle. Antibody was diluted at
1:100(4° overnight). High-pressure and
temperature Tris-EDTA,pH8.0 was used
for antigen retrieval. Negative contrl
(right) obtained from antibody was pre-
absorbed by immunogen peptide.

1,primary
Antibody(red) was diluted at 1:200(4°C
overnight).

2, Goat Anti Rabbit IgG (H&L) - Alexa
Fluor 594 Secondary antibody was
diluted at 1:1000(room temperature,
50min)

3, Picture
B: DAPI(blue) 10min.