

| | |
|--------------------------|---|
| Cat. No: | MABN81703 |
| Conjugate: | Unconjugated |
| Size: | 100µL |
| Clone: | Monoclonal |
| Concentration: | 1mg/ml |
| Host: | Mouse |
| Isotype: | Mouse IgG2b |
| Immunogen: | Purified recombinant fragment of human VIMP (AA: 1-187) expressed in E. Coli. |
| Reactivity: | Human |
| Applications: | WB 1:500-1:2000,IHC 1:200-1:1000,ELISA 1:5000-1:20000 |
| Molecular Weight: | 21.2kDa |
| Purification: | Affinity Purification |
| Synonyms: | SELS; ADO15; SBB18; SEPS1; AD-015 |

Background: This gene encodes a member of the selenoprotein family, characterized by a selenocysteine (Sec) residue at the active site. The selenocysteine is encoded by the UGA codon that normally signals translation termination. The 3' UTR of selenoprotein genes have a common stem-loop structure, the sec insertion sequence (SECIS), that is necessary for the recognition of UGA as a Sec codon rather than as a stop signal. Studies suggest that this protein may regulate cytokine production, and thus play a key role in the control of the inflammatory response. Alternative splicing results in multiple transcript variants encoding different isoforms.

| | |
|-----------------|--|
| Form: | Liquid |
| Buffer: | Purified antibody in PBS with 0.05% sodium azide |
| Storage: | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles. |