

Cat. No:	MABN83814
Conjugate:	Unconjugated
Size:	100µL
Clone:	Monoclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	A synthesized peptide derived from human WASF2
Reactivity:	Human
Applications:	WB 1:1000-1:2000,IHC 1:100-1:200,ICC/IF 1:50-1:200,ICC 1:50-1:200
Molecular Weight:	Calculated MW: 54 kDa ; Observed MW: 80 kDa
Purification:	Affinity Purification
Synonyms:	Wiskott-Aldrich syndrome protein family member 2WASP family protein member 2; Protein WAVE-2; Verprolin homology domain-containing protein 2; WASF2;;WASF2
Background:	Downstream effector molecule involved in the transmission of signals from tyrosine kinase receptors and small GTPases to the actin cytoskeleton. Promotes formation of actin filaments. Part of the WAVE complex that regulates lamellipodia formation. The WAVE complex regulates actin filament reorganization via its interaction with the Arp2/3 complex.
Form:	Liquid
Buffer:	Purified antibody in PBS with 0.05% sodium azide,0.05% BSA and 50% glycerol.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only
IMMUNOLOGICAL SCIENCES**