

Cat. No:	MABN02770
Conjugate:	Unconjugated
Size:	100µL
Clone:	Monoclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	Recombinant protein of human WAVE 1
Reactivity:	Human,Mouse,Rat
Applications:	WB 1:500-1:1000
Molecular Weight:	Calculated MW: 62 kDa; Observed MW: 90 kDa
Purification:	Affinity Purification
Synonyms:	WAVE; SCAR1; WAVE1
Background:	<p>Downstream effector molecule involved in the transmission of signals from tyrosine kinase receptors and small GTPases to the actin cytoskeleton. Promotes formation of actin filaments. Part of the WAVE complex that regulates lamellipodia formation. The WAVE complex regulates actin filament reorganization via its interaction with the Arp2/3 complex . As component of the WAVE1 complex, required for BDNF-NTRK2 endocytic trafficking and signaling from early endosomes .</p>
Form:	Liquid
Buffer:	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only
IMMUNOLOGICAL SCIENCES**