

<b>Cat. No:</b>	MABN03257
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Monoclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	A synthetic peptide of human XRCC1
<b>Reactivity:</b>	Human,Rat
<b>Applications:</b>	WB 1:500-1:1000,ICC/IF 1:50-1:200
<b>Molecular Weight:</b>	Calculated MW: 69 kDa; Observed MW: 82 kDa
<b>Purification:</b>	Affinity Purification
<b>Synonyms:</b>	XRCC1; DNA repair protein XRCC1; X-ray repair cross-complementing protein 1
<b>Background:</b>	XRCC1 acts as a scaffold protein to coordinate DNA abasic site repair through interaction with several other repair proteins. At least eight XRCC1 protein partners have been identified, including the polynucleotide kinase PNK, DNA ligase III, poly (ADP-ribose) polymerase, and PCNA.
<b>Form:</b>	Liquid
<b>Buffer:</b>	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only  
IMMUNOLOGICAL SCIENCES**