

<b>Cat. No:</b>	MABN80819
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Monoclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Mouse
<b>Isotype:</b>	Mouse IgG1
<b>Immunogen:</b>	Purified recombinant fragment of human ZBTB7B expressed in E. Coli.
<b>Reactivity:</b>	Human
<b>Applications:</b>	WB 1:500-1:2000,IHC 1:200-1:1000,ICC 1:200-1:1000,ELISA 1:5000-1:20000
<b>Molecular Weight:</b>	58kDa
<b>Purification:</b>	Affinity Purification
<b>Synonyms:</b>	THPOK; ZFP67; ZBTB15; c-Krox; hcKrox; ZNF857B; DKFZp686G01254; ZBTB7B
<b>Background:</b>	<p>ZBTB7B is a transcription regulator that acts as a key regulator of lineage commitment of immature T-cell precursors. It is necessary and sufficient for commitment of CD4 lineage, while its absence causes CD8 commitment. Development of immature T-cell precursors (thymocytes) to either the CD4 helper or CD8 killer T-cell lineages correlates precisely with their T-cell receptor specificity for major histocompatibility complex class II or class I molecules, respectively. ZBTB7B is a transcriptional repressor of the collagen COL1A1 and COL1A2 genes. It may also function as a repressor of fibronectin and possibly other extracellular matrix genes.</p>
<b>Form:</b>	Liquid
<b>Buffer:</b>	Ascitic fluid containing 0.03% sodium azide.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only  
IMMUNOLOGICAL SCIENCES**