

<b>Cat. No:</b>	ABN20092
<b>Conjugate:</b>	Unconjugated
<b>Size:</b>	100µL
<b>Clone:</b>	Polyclonal
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rabbit
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	The antiserum was produced against synthesized peptide derived from the Internal region of human ZFP91. AA range:401-450
<b>Reactivity:</b>	Human,Mouse,Rat
<b>Applications:</b>	WB 1:500-1:2000,ELISA 1:10000-1:20000
<b>Molecular Weight:</b>	60kDa
<b>Purification:</b>	Affinity purification
<b>Synonyms:</b>	ZFP91 ZNF757 FKSG11

<b>Background:</b>	<p>The protein encoded by this gene is a member of the zinc finger family of proteins. The gene product contains C2H2-type domains, which are the classical zinc finger domains found in numerous nucleic acid-binding proteins. This protein functions as a regulator of the non-canonical NF-kappaB pathway in lymphotoxin-beta receptor signaling. Alternative splicing results in multiple transcript variants. A read-through transcript variant composed of ZFP91 and the downstream CNTF gene sequence has been identified, but it is thought to be non-coding. Read-through transcription of ZFP91 and CNTF has also been observed in mouse. A ZFP91-related pseudogene has also been identified on chromosome 2. [provided by RefSeq, Oct 2010],disease:Overexpressed in most acute myelogenous leukemia (AML) cases (27 over 29).,function:CNTF is a survival factor for various neuronal cell types. Seems to prevent the degeneration of motor axons after axotomy.,function:May be involved in transcriptional regulation. May play an important role in cell proliferation and/or anti-apoptosis.,online information:Ciliary neurotrophic factor entry,similarity:Belongs to the CNTF family.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 5 C2H2-type zinc fingers.,subunit:Homodimer.,tissue specificity:Expressed ubiquitously, particularly at high level in testis. Isoform 2 is testis specific.,tissue specificity:Nervous system.,</p>
<b>Form:</b>	Liquid
<b>Buffer:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Storage:</b>	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

**For Research use only  
IMMUNOLOGICAL SCIENCES**