



<b>Cat. No:</b>	AB-84194
<b>Size:</b>	100 ug
<b>Clone:</b>	POLY
<b>Concentration:</b>	1mg/ml
<b>Host:</b>	Rb
<b>Isotype:</b>	IgG
<b>Immunogen:</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 676-895 of human ZNF281.
<b>Reactivity:</b>	Hu, Ms, Rt
<b>Applications:</b>	Western Blot: 1:500 - 1:2000 Immunohistochemistry: 1:50 - 1:200
<b>Molecular Weight:</b>	110kDa
<b>Purification:</b>	Aff. Pur.
<b>Synonyms:</b>	ZNF281; ZBP-99; ZNP-99; zinc finger protein 281.
<b>Form:</b>	Liquid
<b>Buffer:</b>	PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.
<b>Storage:</b>	Store at -20°C. Avoid freeze / thaw cycles.



Western blot analysis of extracts of mouse spleen, using ZNF281 antibody at 1:3000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.  
Lysates/proteins: 25ug per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL West Pico Plus.  
Exposure time: 10s.



Western blot analysis of lysates from Hela, A549, MCF-7 cell line, mouse liver, mouse small intestine tissue lysate (from left to right), using ZNF281 Antibody (Center) ZNF281 was diluted at 1: 1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1: 10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

**For Research use only  
IMMUNOLOGICAL SCIENCES**