

Product name:	3pK Rabbit Polyclonal Antibody
Cat number:	ABN06320
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human MAPK3. AA range:301-350
Reactivity:	Human,Mouse,Rat
Applications:	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000
Molecular Weight:	42kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

This gene encodes a member of the Ser/Thr protein kinase family. This kinase functions as a mitogen-activated protein kinase (MAP kinase)- activated protein kinase. MAP kinases are also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This kinase was shown to be activated by growth inducers and stress stimulation of cells. In vitro studies demonstrated that ERK, p38 MAP kinase and Jun N-terminal kinase were all able to phosphorylate and activate this kinase, which suggested the role of this kinase as an integrative element of signaling in both mitogen and stress responses. This kinase was reported to interact with, phosphorylate and repress the activity of E47, which is a basic helix-loop-helix transcription factor known to be involved in the regulation of tissue-specific gene expression and catalytic activity: $\text{ATP} + \text{a protein} = \text{ADP} + \text{a phosphoprotein}$., function: Modulator of polycomb-mediated repression, which can be activated either by ERK, p38 and JNK. Substrate of CSBP. In vitro, phosphorylates HSPB1, BMI1/PCGF4 and TCF3., similarity: Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family., similarity: Contains 1 protein kinase domain., subcellular location: Predominantly located in the nucleus, when activated it translocates to the cytoplasm., subunit: Interacts with TCF3 and with polycomb proteins, such as PCH2 and BMI1/PCGF4., tissue specificity: Widely expressed, with a higher expression level observed in heart and skeletal muscle. No expression in brain.,