

|                          |                                                                                                                                          |
|--------------------------|------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Product name:</b>     | 4E-BP1 (phospho Thr37) Rabbit Polyclonal Antibody                                                                                        |
| <b>Cat number:</b>       | ABN04183                                                                                                                                 |
| <b>Conjugate:</b>        | Unconjugated                                                                                                                             |
| <b>Size:</b>             | 100µL                                                                                                                                    |
| <b>Clone:</b>            | Polyclonal                                                                                                                               |
| <b>Concentration:</b>    | 1mg/ml                                                                                                                                   |
| <b>Host:</b>             | Rabbit                                                                                                                                   |
| <b>Isotype:</b>          | IgG                                                                                                                                      |
| <b>Immunogen:</b>        | The antiserum was produced against synthesized peptide derived from human 4E-BP1 around the phosphorylation site of Thr36. AA range:4-53 |
| <b>Reactivity:</b>       | Human,Mouse,Rat,Rabbit,Fish,Chicken,Guinea pig,Sheep,Bovine                                                                              |
| <b>Applications:</b>     | WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000                                                                   |
| <b>Molecular Weight:</b> | 18kDa                                                                                                                                    |
| <b>Purification:</b>     | Affinity purification                                                                                                                    |
| <b>Form:</b>             | Liquid                                                                                                                                   |
| <b>Buffer:</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.                                                       |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.                                             |

**Background:**

eukaryotic translation initiation factor 4E binding protein 1(EIF4EBP1) Homo sapiens This gene encodes one member of a family of translation repressor proteins. The protein directly interacts with eukaryotic translation initiation factor 4E (eIF4E), which is a limiting component of the multisubunit complex that recruits 40S ribosomal subunits to the 5' end of mRNAs. Interaction of this protein with eIF4E inhibits complex assembly and represses translation. This protein is phosphorylated in response to various signals including UV irradiation and insulin signaling, resulting in its dissociation from eIF4E and activation of mRNA translation. [provided by RefSeq, Jul 2008],function:Regulates eIF4E activity by preventing its assembly into the eIF4F complex. Mediates the regulation of protein translation by hormones, growth factors and other stimuli that signal through the MAP kinase pathway.,PTM:Phosphorylated on serine and threonine residues in response to insulin, EGF and PDGF. Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the eIF4E-binding protein family.,subunit:Nonphosphorylated EIF4EBP1 competes with EIF4G1/EIF4G3 to interact with EIF4E; insulin stimulated MAP-kinase (MAPK1 and MAPK3) phosphorylation of EIF4EBP1 causes dissociation of the complex allowing EIF4G1/EIF4G3 to bind and consequent initiation of translation. Rapamycin can attenuate insulin stimulation, mediated by FKBP.,