

Product name:	α -tubulin(Acetyl Lys40)(5H5)Mouse Monoclonal Antibody
Cat number:	MABN06271
Conjugate:	Unconjugated
Size:	100 μ L
Clone:	Monoclonal
Concentration:	1mg/ml
Host:	Mouse
Isotype:	IgG
Immunogen:	Synthetic Peptide of α -tubulin (Acetyl Lys40)
Reactivity:	Human,Rat,Mouse
Applications:	WB 1:1000-1:2000,IHC 1:100-1:200,ICC/IF 1:50-1:200
Molecular Weight:	52kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	function:Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha-chain.,PTM:Undergoes a tyrosination/detyrosination cycle, the cyclic removal and re-addition of a C-terminal tyrosine residue by the enzymes tubulin tyrosine carboxypeptidase (TTCP) and tubulin tyrosine ligase (TTL), respectively.,similarity:Belongs to the tubulin family.,subunit:Dimer of alpha and beta chains.,function:Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha-chain.,PTM:Undergoes a tyrosination/detyrosination cycle, the cyclic removal and re-addition of a C-terminal tyrosine residue by the enzymes tubulin tyrosine carboxypeptidase (TTCP) and tubulin tyrosine ligase (TTL), respectively.,similarity:Belongs to the tubulin family.,subunit:Dimer of alpha and beta chains.,