

Product name:	ADAMTS-19 Rabbit Polyclonal Antibody
Cat number:	ABN06600
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human ADAMTS19. AA range:251-300
Reactivity:	Human,Mouse
Applications:	WB 1:500-1:2000,ELISA 1:20000-1:40000
Molecular Weight:	140kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	This gene encodes a member of the ADAMTS (a disintegrin and metalloproteinase with thrombospondin motif) protein family. Members of the family share several distinct protein modules, including a propeptide region, a metalloproteinase domain, a disintegrin-like domain, and a thrombospondin type 1 (TS) motif. Individual members of this family differ in the number of C-terminal TS motifs, and some have unique C-terminal domains. The protein encoded by this gene has high sequence similarity to the protein encoded by ADAMTS16, another family member. [provided by RefSeq, Jul 2008],cofactor: Binds 1 zinc ion per subunit.,domain: The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.,PTM: The precursor is cleaved by a furin endopeptidase.,similarity: Contains 1 disintegrin domain.,similarity: Contains 1 peptidase M12B domain.,similarity: Contains 1 PLAC domain.,similarity: Contains 5 TSP type-1 domains.,tissue specificity: Expressed in fetal lung, but not in any adult tissues examined. Expression was detected in an osteosarcoma cDNA library.,