






Product name:	Anti-Ki67 Rabbit pAb
Cat number:	ABS111141
Conjugate:	unconjugated
Size:	100 µL
Clone:	Poly
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	KLH conjugated Synthetic peptide corresponding to Mouse Ki67
Reactivity:	Mouse, Rat
Applications:	IHC/IF 1: 500-1: 1200
Purification:	Affinity purification
Buffer:	PBS with 0.02% sodium azide, 100 µg/ml BSA and 50% glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/ thaw cycles.




IHC analysis of ki67 (GB111141).
Sample: Mouse placenta (Paraffin), 4% PFA (G1101) 12-24h.
Antigen retrieval: Citrate buffer (pH 6.0) (G1201), pressure cooker 2min.
Blocking buffer: 3% BSA in PBS (GC305010), RT, 30min.
Primary antibody: 1: 1000, 4°C overnight.
Secondary antibody: HRP Goat Anti-Rabbit IgG (GB23303), 1: 200 RT 1h.




IHC analysis of Ki67 (GB111141).
Sample: Mouse subcutaneous transplant tumor[H22] (Paraffin), 4% PFA (G1101) 12-24h.
Antigen retrieval: Citrate buffer (pH 6.0) (G1201), pressure cooker 2min.
Blocking buffer: 3% BSA in PBS (GC305010), RT, 30min.
Primary antibody: 1: 500, 4°C overnight.
Secondary antibody: HRP Goat Anti-Rabbit IgG (GB23303), 1: 200 RT 1h.



IHC analysis of ki67 (GB111141).
Sample: Rat colon (Paraffin), 4% PFA (G1101) 12-24h.
Antigen retrieval: Citrate buffer (pH 6.0) (G1201), pressure cooker 2min.
Blocking buffer: 3% BSA in PBS (GC305010), RT, 30min.
Primary antibody: 1: 750, 4°C overnight.
Secondary antibody: HRP Goat Anti-Rabbit IgG (GB23303), 1: 200 RT 1h.



IF analysis of ki67 (GB111141).
Sample: Mouse testis (Paraffin), 4% PFA (G1101) 12-24h.
Antigen retrieval: TE buffer (pH 9.0) (G1203), 98°C, 20min.
Blocking buffer: 3% BSA in PBS (GC305010), RT, 30min.
Primary antibody: 1: 600, 4°C overnight.
Secondary antibody: Cy3 conjugated Goat Anti-Rabbit IgG (H+L) (GB21303), 1: 300 RT, 1h.



IF analysis of ki67 (GB111141).
Sample: Rat colon (Paraffin), 4% PFA (G1101) 12-24h.
Antigen retrieval: TE buffer (pH 9.0) (G1203), 98°C, 20min.
Blocking buffer: 3% BSA in PBS (GC305010), RT, 30min.
Primary antibody: 1: 600, 4°C overnight.
Secondary antibody: Cy3 conjugated Goat Anti-Rabbit IgG (H+L) (GB21303), 1: 300 RT, 1h.