

|                          |                                                                                                |
|--------------------------|------------------------------------------------------------------------------------------------|
| <b>Product name:</b>     | ATF-2 Rabbit Polyclonal Antibody                                                               |
| <b>Cat number:</b>       | ABN07265                                                                                       |
| <b>Conjugate:</b>        | Unconjugated                                                                                   |
| <b>Size:</b>             | 100µL                                                                                          |
| <b>Clone:</b>            | Polyclonal                                                                                     |
| <b>Concentration:</b>    | 1mg/ml                                                                                         |
| <b>Host:</b>             | Rabbit                                                                                         |
| <b>Isotype:</b>          | IgG                                                                                            |
| <b>Immunogen:</b>        | The antiserum was produced against synthesized peptide derived from human ATF2. AA range:40-89 |
| <b>Reactivity:</b>       | Human,Mouse,Rat                                                                                |
| <b>Applications:</b>     | WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000,IP 1:20-1:50           |
| <b>Molecular Weight:</b> | 52kDa                                                                                          |
| <b>Purification:</b>     | Affinity purification                                                                          |
| <b>Form:</b>             | Liquid                                                                                         |
| <b>Buffer:</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.             |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.   |

**Background:**

activating transcription factor 2(ATF2) Homo sapiens This gene encodes a transcription factor that is a member of the leucine zipper family of DNA binding proteins. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions This protein binds to the cAMP-responsive element (CRE), an octameric palindrome. It forms a homodimer or a heterodimer with c-Jun and stimulates CRE-dependent transcription. This protein is also a histone acetyltransferase (HAT) that specifically acetylates histones H2B and H4 in vitro; thus it may represent a class of sequence-specific factors that activate transcription by direct effects on chromatin components. The encoded protein may also be involved in cell's DNA damage response independent of its role in transcriptional regulation. Several alternatively spliced transcript variants have been found for this gene [provided by RefSeq, Jan 2014]caution:It is uncertain whether Met-1 or Met-19 is the initiator.,function:Transcriptional activator, probably constitutive, which binds to the cAMP-responsive element (CRE) (consensus: 5'-GTGACGT[AC][AG]-3'), a sequence present in many viral and cellular promoters. Interaction with JUN redirects JUN to bind to CREs preferentially over the 12-O-tetradecanoylphorbol-13-acetate response elements (TREs) as part of an ATF2-c-Jun complex.,PTM:Phosphorylation of Thr-69 and Thr-71 by MAPK14 causes increased transcriptional activity. Also phosphorylated and activated by JNK.,similarity:Belongs to the bZIP family.,similarity:Belongs to the bZIP family. ATF subfamily.,similarity:Contains 1 bZIP domain.,similarity:Contains 1 C2H2-type zinc finger.,subunit:Binds DNA as a dimer and can form a homodimer in the absence of DNA. Can form a heterodimer with JUN. Interacts with SMAD3 and SMAD4. Binds through its N-terminal region to UTF1 which acts as a coactivator of ATF2 transcriptional activity.,tissue specificity:Abundant expression seen in the brain.,