

Product name:	ATP5I Rabbit Polyclonal Antibody
Cat number:	ABN07335
Conjugate:	Unconjugated
Size:	100μL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human ATP5I. AA range:20-69
Reactivity:	Human,Rat,Mouse
Applications:	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
Molecular Weight:	8kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner membrane during oxidative phosphorylation. It is composed of two linked multi-subunit complexes: the soluble catalytic core, F₁, and the membrane-spanning component, F_o, which comprises the proton channel. The F₁ complex consists of 5 different subunits (alpha, beta, gamma, delta, and epsilon) assembled in a ratio of 3 alpha, 3 beta, and a single representative of the other 3. The F_o seems to have nine subunits (a, b, c, d, e, f, g, F₆ and 8). This gene encodes the e subunit of the F_o complex. Alternative splicing results in multiple transcript variants.[provided by RefSeq, Jun 2010],function:Mitochondrial membrane ATP synthase (F₁)F_o ATP synthase or Complex V) produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F₁ - containing the extramembraneous catalytic core, and F_o - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F₁ is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Part of the complex F_o domain. Minor subunit located with subunit a in the membrane.,similarity:Belongs to the ATPase e subunit family.,subunit:F-type ATPases have 2 components, CF₁ - the catalytic core - and CF_o - the membrane proton channel. CF_o seems to have nine subunits: a, b, c, d, e, f, g, F₆ and 8 (or A6L).,