

Product name:	BCL-2
Cat number:	AB-10227
Conjugate:	Unconjugated
Size:	100 ug
Clone:	POLY
Concentration:	1mg/ml
Host:	Rb
Isotype:	IgG
Immunogen:	Peptide derived from the N-terminal sequence of human bcl2. Antibody recognizes the epitope located between Ala42 - Ala60.
Reactivity:	Hu
Applications:	WB: 1:5,000 ELISA - 1:50,000 - 1:100,000 ICC - 1:100 - 1:300 IP: 1:100 - 1:300
Purification:	Aff. Pur.
Form:	Liquid
Buffer:	20 mM Tris-HCl, pH 8.0 , 10 mg/ml BSA, 0.05% Sodium Azide
Storage:	10 µl aliquots at -20°C. Avoid repeated freezing and thawing
Background:	Bcl-2 family of proteins is a key regulator of apoptosis that function to either inhibit or promote cell death. The over expression of members such as Bcl-2 and Bcl-xL inhibit the apoptotic process (1,2). The Bcl-2 family members are also characterized by dimerizing to further modulate apoptosis. Bag-1, for example, has been found to form a heterodimer with Bcl-2 resulting in the enhancement of the anti-apoptotic effect of Bcl-2 (3,4). Other anti-apoptotic Bcl-2 family members include A1, Bcl-xg, Bcl-xb, Mcl-1, BAR, BI-1 and Bcl-w (5). The pro-apoptotic family members include Bax, Bcl-xS, Bad, Bak, NBK, BID, Hrk, Bok, Bim, Noxa and Diva. Bax and Bak have been shown to play a critical role in cytochrome c release from mitochondria and thus initiate apoptosis (6). Bad plays a critical role in the Bax-mediated apoptosis pathway by dimerizing with Bcl-xL, causing the displacement of Bax. The displacement of Bax allows apoptosis to proceed (7). Bcl-xS, a shorter version of Bcl-xL (lacking amino acids 126-188), apparently utilizes a different pathway than Bax to induce cell death. Some research suggests that Bcl-xS uses a novel mechanism for regulating caspase or it may use an alternate cell death effector pathway (8,9).