

Product name:	CAD Rabbit Polyclonal Antibody
Cat number:	ABN07823
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human CAD. AA range:422-471
Reactivity:	Human,Mouse
Applications:	IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
Molecular Weight:	250kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

The de novo synthesis of pyrimidine nucleotides is required for mammalian cells to proliferate. This gene encodes a trifunctional protein which is associated with the enzymatic activities of the first 3 enzymes in the 6-step pathway of pyrimidine biosynthesis: carbamoylphosphate synthetase (CPS II), aspartate transcarbamoylase, and dihydroorotase. This protein is regulated by the mitogen-activated protein kinase (MAPK) cascade, which indicates a direct link between activation of the MAPK cascade and de novo biosynthesis of pyrimidine nucleotides. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Apr 2015], catalytic activity: (S)-dihydroorotate + H₂O = N-carbamoyl-L-aspartate., catalytic activity: 2 ATP + L-glutamine + HCO₃⁻ + H₂O = 2 ADP + phosphate + L-glutamate + carbamoyl phosphate., catalytic activity: Carbamoyl phosphate + L-aspartate = phosphate + N-carbamoyl-L-aspartate., cofactor: Binds 1 zinc ion per subunit (for dihydroorotase activity) ., enzyme regulation: Allosterically regulated and controlled by phosphorylation. 5-phosphoribose 1-diphosphate is an activator while UMP is an inhibitor of the CPSase reaction., function: This protein is a "fusion" protein encoding four enzymatic activities of the pyrimidine pathway (GATase, CPSase, ATCase and DHOase)., miscellaneous: GATase (glutamine amidotransferase) and CPSase (carbamoyl phosphate synthase) form together the glutamine-dependent CPSase (GD-CPSase) (EC 6.3.5.5)., online information: Aspartate carbamoyltransferase entry, pathway: Pyrimidine metabolism; UMP biosynthesis via de novo pathway; UMP from HCO₃⁻: step 1/6., pathway: Pyrimidine metabolism; UMP biosynthesis via de novo pathway; UMP from HCO₃⁻: step 2/6., pathway: Pyrimidine metabolism; UMP biosynthesis via de novo pathway; UMP from HCO₃⁻: step 3/6., similarity: Belongs to the ATCase/OTCase family., similarity: Contains 1 glutamine amidotransferase type-1 domain., similarity: Contains 2 ATP-grasp domains., similarity: In the central section; belongs to the DHOase family., subunit: Homohexamer.,