



Product name:	Catenin gamma / CTNNG1
Cat number:	MABX-80096
Size:	100 ug
Clone:	CN-3
Concentration:	1mg/ml
Host:	Ms
Isotype:	IgG1
Immunogen:	Recombinant chicken plakoglobin.
Reactivity:	Hu, Rt, Ch
Applications:	Western blot: 2µg/ml Immunocytochemistry: 1µg/ml Immunohistochemistry(Frozen Section): 4µg/ml
Purification:	Purified
Form:	Liquid
Buffer:	Mouse ascites fluid, 1.2% sodium acetate, 2mg BSA, with 0.01mg NaN ₃ as preservative.
Storage:	At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquoted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.
Background:	Junction Plakoglobin(JUP) ,also knows as catenin gamma, is a major cytoplasmic protein that occurs in a soluble and a membrane-associated form and is the only known constituent common to the submembranous plaques of both kinds of adhering junctions, the desmosomes and the intermediate junctions. It is a component of the cadherin-catenin complex, which is predominantly localized where actin filaments anchor in adherens junctions of epithelial cells. The human plakoglobin gene localizes on chromosome 17q21. Gamma-catenin is regulated by the APC tumor suppressor and its oncogenic activity is distinct from that of beta-catenin.



Anti-Catenin gamma(Plakoglobin) antibody (monoclonal),
Western blotting Lane 1: MCF-7 Cell Lysate Lane 2: HELA Cell
Lysate Lane 3: MM231 Cell Lysate Lane 4: HT1080 Cell Lysate



Immunocytochemistry/Immunofluorescence: gamma Catenin Antibody Staining gamma Catenin in A431 cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.