

Product name:	CD40L (15O10) Rabbit Monoclonal Antibody
Cat number:	MABN08392
Conjugate:	Unconjugated
Size:	100µL
Clone:	Monoclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	A synthetic peptide of human CD40L
Reactivity:	Human, Mouse
Applications:	WB 1:1000-1:5000
Molecular Weight:	29kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	Mediates B-cell proliferation in the absence of co-stimulus as well as IgE production in the presence of IL-4. Involved in immunoglobulin class switching. Cytokine that acts as a ligand to CD40/TNFRSF5 (PubMed:1280226, PubMed:31331973). Costimulates T-cell proliferation and cytokine production (PubMed:8617933). Its cross-linking on T-cells generates a costimulatory signal which enhances the production of IL4 and IL10 in conjunction with the TCR/CD3 ligation and CD28 costimulation (PubMed:8617933). Induces the activation of NF-kappa-B (PubMed:15067037, PubMed:31331973). Induces the activation of kinases MAPK8 and PAK2 in T-cells (PubMed:15067037). Induces tyrosine phosphorylation of isoform 3 of CD28 (PubMed:15067037). Mediates B-cell proliferation in the absence of co-stimulus as well as IgE production in the presence of IL4 (By similarity). Involved in immunoglobulin class switching (By similarity).