

Product name:	CD89 Rabbit Polyclonal Antibody
Cat number:	ABN08480
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from the Internal region of human FCAR. AA range:31-80
Reactivity:	Human,Rat,Mouse
Applications:	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
Molecular Weight:	32kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	This gene is a member of the immunoglobulin gene superfamily and encodes a receptor for the Fc region of IgA. The receptor is a transmembrane glycoprotein present on the surface of myeloid lineage cells such as neutrophils, monocytes, macrophages, and eosinophils, where it mediates immunologic responses to pathogens. It interacts with IgA-opsonized targets and triggers several immunologic defense processes, including phagocytosis, antibody-dependent cell-mediated cytotoxicity, and stimulation of the release of inflammatory mediators. Multiple alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Jul 2008],alternative products:Additional isoforms seem to exist,function:Binds to the Fc region of immunoglobulins alpha. Mediates several functions including cytokine production.,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like) domains.,subunit:Associates with the Fc epsilon RI gamma 2 receptor inducing tyrosine phosphorylation of gamma 2.,tissue specificity:Differentially expressed between blood and mucosal myeloid cells. Monocytes express isoform A.1, isoform A.2 and isoform A.3 while alveolar macrophages express isoform A.1 and isoform A.2; however they express only one isoform at their surfaces.,