

Product name:	CdcA7 Rabbit Polyclonal Antibody
Cat number:	ABN08535
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human CDCA7. AA range:141-190
Reactivity:	Human,Mouse,Rat
Applications:	WB 1:500-1:2000,ELISA 1:10000-1:20000
Molecular Weight:	43kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	cell division cycle associated 7(CDCA7) Homo sapiens This gene was identified as a c-Myc responsive gene, and behaves as a direct c-Myc target gene. Overexpression of this gene is found to enhance the transformation of lymphoblastoid cells, and it complements a transformation-defective Myc Box II mutant, suggesting its involvement in c-Myc-mediated cell transformation. Two alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008],function:Participates in MYC-mediated cell transformation; induces anchorage-independent growth and clonogenicity in lymphoblastoid cells. Insufficient to induce tumorigenicity when overexpressed but contributes to MYC-mediated tumorigenesis. May play a role as transcriptional regulator.,induction:Activated by MYC and possibly E2F1.,miscellaneous:CDCA7 expression is correlated with MYC expression in lymphoblastoid, lymphoma and breast cancer cell lines.,tissue specificity:Ubiquitous with higher level in thymus and small intestine. Overexpressed in a large number of tumors, in blood from patients with acute myelogenous leukemia (AML) and in chronic myelogenous leukemia (CML) blast crisis.,