




Product name:	Cleaved-Caspase-3 p17 (D175) Rabbit Polyclonal Antibody
Cat number:	ABE1002
Conjugate:	unconjugated
Size:	100 ug
Clone:	Polyclonal
Concentration:	1 mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human Caspase 3. AA range:126-175
Reactivity:	Human;Mouse;Rat
Applications:	WB 1:500-2000, IHC-p 1:50-300, IF 1:50-300
Molecular Weight:	17-34kD
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Form:	liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5%BSAand0.02% sodium azide.
Storage:	-20°C/1 year
Background:	This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein cleaves and activates caspases 6, 7 and 9, and the protein itself is processed by caspases 8, 9 and 10. It is the predominant caspase involved in the cleavage of amyloid-beta 4A precursor protein, which is associated with neuronal death in Alzheimer's disease. Alternative splicing of this gene results in two transcript variants that encode the same protein. [provided by RefSeq, Jul 2008],




Western Blot analysis of 293 cells using Cleaved-Caspase-3 p17 (D175) Polyclonal Antibody




Post-surgical wound management and prevention of triple-negative breast cancer recurrence with a pyroptosis-inducing, photopolymerizable hydrogel
JOURNAL OF CONTROLLED RELEASE Sanjun Shi WB
Mouse 4 T1 cell




Zhou, Li, et al. "Coptisine induces apoptosis in human hepatoma cells through activating 67-kDa laminin receptor/cGMP signaling." *Frontiers in pharmacology* 9 (2018).




Bu, Xuefeng, et al. "Recombinant Newcastle disease virus (rL-RVG) triggers autophagy and apoptosis in gastric carcinoma cells by inducing ER stress." *American journal of cancer research* 6.5 (2016): 924.




Negative Control
Immunohistochemical analysis of paraffin-embedded Human-colon tissue. 1. Cleaved-Caspase-3 p17 (D175) Polyclonal Antibody was diluted at 1:200(4°C, overnight). 2. Sodium citrate pH 6.0 was used for antibody retrieval(>98°C, 20min). 3. Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.




Negative Control
Immunohistochemical analysis of paraffin-embedded Rat-heart tissue. 1. Cleaved-Caspase-3 p17 (D175) Polyclonal Antibody was diluted at 1:200(4°C, overnight). 2. Sodium citrate pH 6.0 was used for antibody retrieval(>98°C, 20min). 3. Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.



Negative Control
Immunohistochemical analysis of paraffin-embedded Mouse-kidney tissue. 1. Cleaved-Caspase-3 p17 (D175) Polyclonal Antibody was diluted at 1:200(4°C, overnight). 2. Sodium citrate pH 6.0 was used for antibody retrieval(>98°C, 20min). 3. Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.



Immunofluorescence analysis of Human-lung-cancer tissue. 1. Cleaved-Caspase-3 p17 (D175) Polyclonal Antibody(red) was diluted at 1:200(4°C, overnight). 2. Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min). 3. Picture B: DAPI(blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B



Immunofluorescence analysis of Mouse-liver tissue. 1. Cleaved-Caspase-3 p17 (D175) Polyclonal Antibody(red) was diluted at 1:200(4°C, overnight). 2. Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min). 3. Picture B: DAPI(blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B