

Product name:	Cleaved-Plasma Kallikrein HC (R390) Rabbit Polyclonal Antibody
Cat number:	ABN09027
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human KLKB1. AA range:341-390
Reactivity:	Human,Rat,Mouse
Applications:	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
Molecular Weight:	45kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

This gene encodes a glycoprotein that participates in the surface-dependent activation of blood coagulation, fibrinolysis, kinin generation and inflammation. The encoded preproprotein present in plasma as a non-covalent complex with high molecular weight kininogen undergoes proteolytic processing mediated by activated coagulation factor XII to generate a disulfide-linked, heterodimeric serine protease comprised of heavy and light chains. Certain mutations in this gene cause prekallikrein deficiency. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jan 2016], catalytic activity: Cleaves selectively Arg-|-Xaa and Lys-|-Xaa bonds, including Lys-|-Arg and Arg-|-Ser bonds in (human) kininogen to release bradykinin., disease: Defects in KLKB1 are the cause of prekallikrein deficiency (PKK deficiency) [MIM:612423]; also called Fletcher factor deficiency. This disorder is a blood coagulation defect., function: The enzyme cleaves Lys-Arg and Arg-Ser bonds. It activates, in a reciprocal reaction, factor XII after its binding to a negatively charged surface. It also releases bradykinin from HMW kininogen and may also play a role in the renin-angiotensin system by converting prorenin into renin., similarity: Belongs to the peptidase S1 family., similarity: Belongs to the peptidase S1 family. Plasma kallikrein subfamily., similarity: Contains 1 peptidase S1 domain., similarity: Contains 4 apple domains., subunit: The zymogen is activated by factor XIIa, which cleaves the molecule into a light chain, which contains the active site, and a heavy chain, which associates with HMW kininogen. These chains are linked by one or more disulfide bonds.,