

Product name:	COL12A1 Rabbit Polyclonal Antibody
Cat number:	ABN09166
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human Collagen XII alpha1. AA range:1481-1530
Reactivity:	Human,Rat,Mouse
Applications:	IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

This gene encodes the alpha chain of type XII collagen, a member of the FACIT (fibril-associated collagens with interrupted triple helices) collagen family. Type XII collagen is a homotrimer found in association with type I collagen, an association that is thought to modify the interactions between collagen I fibrils and the surrounding matrix. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008], alternative products: The final tissue form of collagen XII may contain homotrimers of either isoform 1 or isoform 2 or any combination of isoform 1 and isoform 2, function: Type XII collagen interacts with type I collagen-containing fibrils, the COL1 domain could be associated with the surface of the fibrils, and the COL2 and NC3 domains may be localized in the perifrillar matrix., PTM: Hydroxylation on proline residues within the sequence motif, GXPG, is most likely to be 4-hydroxy as this fits the requirement for 4-hydroxylation in vertebrates., PTM: O-glycosylation of isoform 1; glycosaminoglycan of chondroitin-sulfate type., PTM: The triple-helical tail is stabilized by disulfide bonds at each end., similarity: Belongs to the fibril-associated collagens with interrupted helices (FACIT) family., similarity: Contains 1 TSP N-terminal (TSPN) domain., similarity: Contains 18 fibronectin type-III domains., similarity: Contains 4 VWFA domains., subunit: Trimer of identical chains each containing 190 kDa of non-triple-helical sequences., tissue specificity: Found in collagen I-containing tissues: both isoform 1 and isoform 2 appear in amnion, chorion, skeletal muscle, small intestine, and in cell culture of dermal fibroblasts, keratinocytes and endothelial cells. Only isoform 2 is found in lung, placenta, kidney and a squamous cell carcinoma cell line. Isoform 1 is also present in the corneal epithelial Bowman's membrane (BM) and the interfibrillar matrix of the corneal stroma, but it is not detected in the limbal BM.,