

Product name:	CP4FN Rabbit Polyclonal Antibody
Cat number:	ABN09304
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	Synthesized peptide derived from human protein . at AA range: 440-520
Reactivity:	Human,Rat,Mouse
Applications:	WB 1:500-1:2000,ELISA 1:5000-1:20000
Molecular Weight:	58kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	cytochrome P450 family 4 subfamily F member 22(CYP4F22) Homo sapiens This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This gene is part of a cluster of cytochrome P450 genes on chromosome 19 and encodes an enzyme thought to play a role in the 12(R)-lipoxygenase pathway. Mutations in this gene are the cause of ichthyosis lamellar type 3. [provided by RefSeq, Jul 2008],cofactor:Heme group.,disease:Defects in CYP4F22 are the cause of ichthyosis lamellar type 3 (LI3) [MIM:604777]. LI is a non-bullous ichthyosis, a skin disorder characterized by abnormal cornification of the epidermis. It is one the most severe forms of ichthyoses apparent at birth and persisting throughout life. LI patients are born encased in a tight, shiny, translucent covering called collodion membrane. Over the first weeks of life, the collodion membrane is gradually replaced by generalized large, dark brown, plate-like scales with minimal to no erythroderma. Tautness of facial skin commonly results in ectropion, eclabium and scarring alopecia of the scalp. Common complications are severe heat intolerance and recurrent ear infections.,similarity:Belongs to the cytochrome P450 family.,