

Product name:	CYP2C8 Rabbit Polyclonal Antibody
Cat number:	ABN09654
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	Synthesized peptide derived from the Internal region of human CYP2C8.
Reactivity:	Human,Rat,Mouse
Applications:	WB 1:500-1:2000,ELISA 1:5000-1:20000
Molecular Weight:	55kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by phenobarbital. The enzyme is known to metabolize many xenobiotics, including the anticonvulsive drug mephenytoin, benzo(a)pyrene, 7-ethoxycoumarin, and the anti-cancer drug taxol. This gene is located within a cluster of cytochrome P450 genes on chromosome 10q24. Several transcript variants encoding a few different isoforms have been found for this gene. [provided by RefSeq, Nov 2010], catalytic activity: $\text{RH} + \text{reduced flavoprotein} + \text{O}_2 = \text{ROH} + \text{oxidized flavoprotein} + \text{H}_2\text{O}$, caution: Alternative splicing has been shown to occur but the shorter forms are believed to be non-functional., cofactor: Heme group., function: Cytochromes P450 are a group of heme-thiolate monooxygenases. In liver microsomes, this enzyme is involved in an NADPH-dependent electron transport pathway. It oxidizes a variety of structurally unrelated compounds, including steroids, fatty acids, and xenobiotics. In the epoxidation of arachidonic acid it generates only 14,15- and 11,12-cis-epoxyeicosatrienoic acids. It is the principal enzyme responsible for the metabolism the anti-cancer drug paclitaxel (taxol)., induction: By phenobarbital., online information: CYP2C8 alleles, similarity: Belongs to the cytochrome P450 family.,