

|                          |                                                                                                     |
|--------------------------|-----------------------------------------------------------------------------------------------------|
| <b>Product name:</b>     | CysLTR1 Rabbit Polyclonal Antibody                                                                  |
| <b>Cat number:</b>       | ABN09683                                                                                            |
| <b>Conjugate:</b>        | Unconjugated                                                                                        |
| <b>Size:</b>             | 100µL                                                                                               |
| <b>Clone:</b>            | Polyclonal                                                                                          |
| <b>Concentration:</b>    | 1mg/ml                                                                                              |
| <b>Host:</b>             | Rabbit                                                                                              |
| <b>Isotype:</b>          | IgG                                                                                                 |
| <b>Immunogen:</b>        | The antiserum was produced against synthesized peptide derived from human CysLTR1. AA range:131-180 |
| <b>Reactivity:</b>       | Human,Rat,Mouse                                                                                     |
| <b>Applications:</b>     | WB 1:500-1:2000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:10000                                            |
| <b>Molecular Weight:</b> | 38kDa                                                                                               |
| <b>Purification:</b>     | Affinity purification                                                                               |
| <b>Form:</b>             | Liquid                                                                                              |
| <b>Buffer:</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.                  |
| <b>Storage:</b>          | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.        |

**Background:**

This gene encodes a member of the G-protein coupled receptor 1 family. The encoded protein is a receptor for cysteinyl leukotrienes, and is involved in mediating bronchoconstriction via activation of a phosphatidylinositol-calcium second messenger system. Activation of the encoded receptor results in contraction and proliferation of bronchial smooth muscle cells, eosinophil migration, and damage to the mucus layer in the lung. Upregulation of this gene is associated with asthma and dysregulation may also be implicated in cancer. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2013],function:Receptor for cysteinyl leukotrienes mediating bronchoconstriction of individuals with and without asthma. Stimulation by LTD4 results in the contraction and proliferation of smooth muscle, edema, eosinophil migration and damage to the mucus layer in the lung. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system. The rank order of affinities for the leukotrienes is LTD4 >> LTE4 = LTC4 >> LTB4.,miscellaneous:Selective antagonists, such as montelukast (Singulair), zafirlukast (Accolate) and pranlukast (Onon), are used in the treatment of the asthma crisis.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Widely expressed, with highest levels in spleen and peripheral blood leukocytes. Lower expression in several tissues, such as lung (mostly in smooth muscle bundles and alveolar macrophages), placenta, small intestine, pancreas, colon and heart.,