

Product name:	Cystatin C Rabbit Polyclonal Antibody
Cat number:	ABN00136
Conjugate:	Unconjugated
Size:	100μL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	Recombinant protein of human Cystatin C
Reactivity:	Human, Mouse, Rat
Applications:	WB 1:500-1:1000, IHC 1:50-1:100
Molecular Weight:	Calculated MW: 16 kDa; Observed MW: 16 kDa
Purification:	Affinity Chromatography
Form:	Liquid
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	Cystatin C is a 14 kDa member of the Cystatin superfamily of cysteine protease inhibitors. Most cell types secrete Cystatin C. Cystatin C inhibits cathepsins, and thereby may function as a tumor suppressor by inhibiting cathepsin mediated tumor cell invasion. In addition, this tumor suppressor function can also be attributed to Cystatin C's ability to antagonize TGF-β1 signaling.