


Product name:	DDDDK-Tag Rabbit Polyclonal Antibody
Cat number:	AB-84802
Conjugate:	Unconjugated
Size:	100 ug
Clone:	POLY
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	DDDDK synthetic peptide conjugated to KLH.
Reactivity:	Species independent
Applications:	Western Blot: 1/1000 - 1/3000 ELISA:1:5000-20000.
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Background:	The DYKDDDDK (FLAG) peptide has been used extensively as a general tag in expression vectors. This peptide can be expressed and detected with the protein of interest as an amino-terminal or carboxy-terminal fusion. N-terminal FLAG vectors provide an Ek cleavage site for removal of the fusion tag. The FLAG peptide is likely to be located on the surface of a fusion protein because of its hydrophilic nature. As a result, the FLAG peptide is more likely to be accessible to antibodies. A FLAG-tag can be used in many different assays that require recognition by an antibody, such as western blotting, immunocytochemistry, immunoprecipitation, flow cytometry, protein purification, and in the study of protein-protein interactions, cell ultrastructure, and protein localization and so on.



Western Blot analysis using FLAG-tag Polyclonal Antibody against HEK293 cells transfected with vector overexpressing FLAG tag (1) and untransfected (2). Secondary antibody was diluted at 1:20000