

Product name:	Emi1 (16Z12) Rabbit Monoclonal Antibody
Cat number:	MABN10439
Conjugate:	Unconjugated
Size:	100µL
Clone:	Monoclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	Recombinant protein of human Emi1
Reactivity:	Human, Mouse, Rat
Applications:	WB 1:1000-1:5000
Molecular Weight:	50kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

Regulates progression through early mitosis by inhibiting the anaphase promoting complex/cyclosome (APC). Binds to the APC activators CDC20 and FZR1/CDH1 to prevent APC activation. Can also bind directly to the APC to inhibit substrate-binding. Regulator of APC activity during mitotic and meiotic cell cycle (PubMed:17485488, PubMed:17234884, PubMed:17875940, PubMed:23708001, PubMed:23708605, PubMed:16921029). During mitotic cell cycle plays a role as both substrate and inhibitor of APC-FZR1 complex (PubMed:29875408, PubMed:17485488, PubMed:17234884, PubMed:17875940, PubMed:23708001, PubMed:23708605, PubMed:16921029). During G1 phase, plays a role as substrate of APC-FZR1 complex E3 ligase (PubMed:29875408). Then switches as an inhibitor of APC-FZR1 complex during S and G2 leading to cell-cycle commitment (PubMed:29875408). As APC inhibitor, prevents the degradation of APC substrates at multiple levels: by interacting with APC and blocking access of APC substrates to the D-box coreceptor, formed by FZR1 and ANAPC10; by suppressing ubiquitin ligation and chain elongation by APC by preventing the UBE2C and UBE2S activities (PubMed:23708605, PubMed:23708001, PubMed:16921029). Plays a role in genome integrity preservation by coordinating DNA replication with mitosis through APC inhibition in interphase to stabilize CCNA2 and GMNN in order to promote mitosis and prevent rereplication and DNA damage-induced cellular senescence (PubMed:17234884, PubMed:17485488, PubMed:17875940). During oocyte maturation, plays a role in meiosis through inactivation of APC-FZR1 complex. Inhibits APC through RPS6KA2 interaction that increases FBXO5 affinity for CDC20 leading to the metaphase arrest of the second meiotic division before fertilization (By similarity). Controls entry into the first meiotic division through inactivation of APC-FZR1 complex (By similarity). Promotes migration and osteogenic differentiation of mesenchymal stem cells (PubMed:29850565).