

|                       |                                                                                                    |
|-----------------------|----------------------------------------------------------------------------------------------------|
| <b>Product name:</b>  | ErbB-4 Rabbit Polyclonal Antibody                                                                  |
| <b>Cat number:</b>    | ABN10574                                                                                           |
| <b>Conjugate:</b>     | Unconjugated                                                                                       |
| <b>Size:</b>          | 100µL                                                                                              |
| <b>Clone:</b>         | Polyclonal                                                                                         |
| <b>Concentration:</b> | 1mg/ml                                                                                             |
| <b>Host:</b>          | Rabbit                                                                                             |
| <b>Isotype:</b>       | IgG                                                                                                |
| <b>Immunogen:</b>     | The antiserum was produced against synthesized peptide derived from human HER4. AA range:1250-1299 |
| <b>Reactivity:</b>    | Human,Mouse,Rat                                                                                    |
| <b>Applications:</b>  | WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000                          |
| <b>Purification:</b>  | Affinity purification                                                                              |
| <b>Form:</b>          | Liquid                                                                                             |
| <b>Buffer:</b>        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.                 |
| <b>Storage:</b>       | Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.       |

**Background:**

This gene is a member of the Tyr protein kinase family and the epidermal growth factor receptor subfamily. It encodes a single-pass type I membrane protein with multiple cysteine rich domains, a transmembrane domain, a tyrosine kinase domain, a phosphatidylinositol-3 kinase binding site and a PDZ domain binding motif. The protein binds to and is activated by neuregulins and other factors and induces a variety of cellular responses including mitogenesis and differentiation. Multiple proteolytic events allow for the release of a cytoplasmic fragment and an extracellular fragment. Mutations in this gene have been associated with cancer. Alternatively spliced variants which encode different protein isoforms have been described; however, not all variants have been fully characterized. [provided by RefSeq, Jul 2008], catalytic activity: ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate., domain: The WW-binding motifs mediate interaction with WWOX., function: Specifically binds and is activated by neuregulins, NRG-2, NRG-3, heparin-binding EGF-like growth factor, betacellulin and NTAK. Interaction with these factors induces cell differentiation. Not activated by EGF, TGF- $\alpha$ , and amphiregulin., PTM: Isoform JM-A is processed but not isoform JM-B. So, they respectively represent cleavable and non-cleavable forms of the receptor., PTM: Ligand-binding increases phosphorylation on tyrosine residues., similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family. EGF receptor subfamily., similarity: Contains 1 protein kinase domain., subunit: Homodimer or heterodimer with each of the other ERBB receptors (Potential). Interacts with PDZ domains of DLG2, DLG3, DLG4 and the syntrophin SNTB2. Interacts with CBFA2T3, MUC1 and WWOX., tissue specificity: Expressed at highest levels in brain, heart, kidney, in addition to skeletal muscle, parathyroid, cerebellum, pituitary, spleen, testis and breast. Lower levels in thymus, lung, salivary gland, and pancreas. Isoform JM-A and isoform JM-B are expressed in cerebellum, but only the isoform JM-B is expressed in the heart.,