

Product name:	FGFR-4 Rabbit Polyclonal Antibody
Cat number:	ABN10950
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from the Internal region of human FGFR4. AA range:91-140
Reactivity:	Human,Rat,Mouse
Applications:	WB 1:500-1:2000,ELISA 1:5000-1:20000
Molecular Weight:	85kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

The protein encoded by this gene is a member of the fibroblast growth factor receptor family, where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein would consist of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. The genomic organization of this gene, compared to members 1-3, encompasses 18 exons rather than 19 or 20. Although alternative splicing has been observed, there is no evidence that the C-terminal half of the Ig-like catalytic activity: $\text{ATP} + \text{a [protein]-L-tyrosine} = \text{ADP} + \text{a [protein]-L-tyrosine phosphate}$.
function: Receptor for acidic fibroblast growth factor. Does not bind to basic fibroblast growth factor. Binds FGF19.
PTM: Glycosylated (By similarity). Phosphorylated on tyrosine residue (By similarity). Phosphorylation requires the presence of a functional (phosphorylated) FGFR1 and not necessarily by means of FGFR heterodimerization.
similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family.
similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family. Fibroblast growth factor receptor subfamily.
similarity: Contains 1 protein kinase domain.
similarity: Contains 3 Ig-like C2-type (immunoglobulin-like) domains.
subcellular location: Isoform 2 may be secreted.
subunit: Interacts with KLB.
tissue specificity: Expressed in gastrointestinal epithelial cells, pancreas, and gastric and pancreatic cancer cell lines.