

Product name:	Fhit Rabbit Polyclonal Antibody
Cat number:	ABN10957
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human FHIT. AA range:81-130
Reactivity:	Human,Rat,Mouse
Applications:	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000
Molecular Weight:	16kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

This gene, a member of the histidine triad gene family, encodes a diadenosine 5',5'''-P₁,P₃-triphosphate hydrolase involved in purine metabolism. The gene encompasses the common fragile site FRA3B on chromosome 3, where carcinogen-induced damage can lead to translocations and aberrant transcripts of this gene. In fact, aberrant transcripts from this gene have been found in about half of all esophageal, stomach, and colon carcinomas. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Oct 2009],catalytic activity:P(1)-P(3)-bis(5'-adenosyl) triphosphate + H₂O = ADP + AMP.,cofactor:Divalent cations. Magnesium, but manganese and to a lesser extent calcium or cobalt can be substituted; but not zinc, cadmium or nickel.,disease:A chromosomal aberration involving FHIT is observed in early onset bilateral and multifocal clear cell renal carcinoma [MIM:144700]. Translocation t(3;8) (3p14.2).,disease:Associated with digestive tract cancers. Numerous tumor types are found to have aberrant forms of FHIT protein due to deletions in a coding region of chromosome 3p14.2 including the fragile site locus FRA3B.,function:Cleaves A-5'-PPP-5'A to yield AMP and ADP. Possible tumor suppressor for specific tissues.,mass spectrometry: PubMed:15007172,similarity:Contains 1 HIT domain.,subunit:Homodimer.,tissue specificity:Low levels expressed in all tissues tested. Phospho-FHIT observed in liver and kidney, but not in brain and lung. Phospho-FHIT undetected in all tested human tumor cell lines.,