

Product name:	GBDR1 Rabbit Polyclonal Antibody
Cat number:	ABN11324
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human UBAC1. AA range:91-140
Reactivity:	Human,Rat,Mouse
Applications:	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
Molecular Weight:	45kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	function:Non-catalytic subunit of the KPC complex that acts as E3 ubiquitin-protein ligase. Required for poly-ubiquitination and proteasome-mediated degradation of CDKN1B during G1 phase of the cell cycle.,pathway:Protein modification; protein ubiquitination.,similarity:Contains 1 STI1 domain.,similarity:Contains 1 ubiquitin-like domain.,similarity:Contains 2 UBA domains.,subunit:Component of the KPC complex composed of RNF123/KPC1 and UBAC1/KPC2. Interacts with RNF123 via its N-terminal domain. Interacts with the proteasome via its N-terminal domain.,tissue specificity:Ubiquitous.,function:Non-catalytic subunit of the KPC complex that acts as E3 ubiquitin-protein ligase. Required for poly-ubiquitination and proteasome-mediated degradation of CDKN1B during G1 phase of the cell cycle.,pathway:Protein modification; protein ubiquitination.,similarity:Contains 1 STI1 domain.,similarity:Contains 1 ubiquitin-like domain.,similarity:Contains 2 UBA domains.,subunit:Component of the KPC complex composed of RNF123/KPC1 and UBAC1/KPC2. Interacts with RNF123 via its N-terminal domain. Interacts with the proteasome via its N-terminal domain.,tissue specificity:Ubiquitous.,