

Product name:	Glutathione Peroxidase 3/GPx-3 Rabbit Monoclonal Antibody
Cat number:	MABN87419
Conjugate:	Unconjugated
Size:	100μL
Clone:	Monoclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	Recombinant protein of mouse Glutathione Peroxidase 3/GPx-3
Reactivity:	Mouse,Rat
Applications:	WB 1:500-1:2000,IHC 1:200-1:500
Molecular Weight:	Calculated MW:25 kDa; Observed MW:25 kDa
Purification:	Affinity Purification
Form:	Liquid
Buffer:	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.
Background:	The protein encoded by this gene belongs to the glutathione peroxidase family, members of which catalyze the reduction of organic hydroperoxides and hydrogen peroxide (H₂O₂) by glutathione, and thereby protect cells against oxidative damage. Several isozymes of this gene family exist in vertebrates, which vary in cellular location and substrate specificity. This isozyme is secreted and is highly expressed in mouse kidney, which appears to be the major source of the enzyme in plasma. It has a role in mouse organogenesis, and dysregulation of this isozyme has been associated with obesity-related metabolic complications, platelet-dependent thrombosis, colitis-associated carcinoma, and thermosensitive phenotype. This isozyme is also a selenoprotein, containing the rare amino acid selenocysteine (Sec) at its active site. Sec is encoded by the UGA codon, which normally signals translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, designated the Sec insertion sequence (SECIS) element, that is necessary for the recognition of UGA as a Sec codon, rather than as a stop signal. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Aug 2016]