

Product name:	GRB10 (phospho Tyr67) Rabbit Polyclonal Antibody
Cat number:	ABN04743
Conjugate:	Unconjugated
Size:	100µL
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	The antiserum was produced against synthesized peptide derived from human GRB10 around the phosphorylation site of Tyr67. AA range:33-82
Reactivity:	Human,Rat,Mouse
Applications:	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000
Molecular Weight:	67kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

The product of this gene belongs to a small family of adapter proteins that are known to interact with a number of receptor tyrosine kinases and signaling molecules. This gene encodes a growth factor receptor-binding protein that interacts with insulin receptors and insulin-like growth-factor receptors. Overexpression of some isoforms of the encoded protein inhibits tyrosine kinase activity and results in growth suppression. This gene is imprinted in a highly isoform- and tissue-specific manner, with expression observed from the paternal allele in the brain, and from the maternal allele in the placental trophoblasts. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Oct 2010], alternative products: Additional isoforms seem to exist, function: Plays a functional role in insulin and IGF-I signaling. May serve to positively link the insulin and IGF-I receptors to an uncharacterized mitogenic signaling pathway. Interacts with the cytoplasmic domain of the autophosphorylated insulin receptor which is then inhibited. The interaction is mediated by the SH2 domain. Also binds activated platelet-derived growth factor receptor and epidermal growth factor receptor., similarity: Belongs to the GRB7/10/14 family., similarity: Contains 1 PH domain., similarity: Contains 1 Ras-associating domain., similarity: Contains 1 SH2 domain., subunit: Interacts with GIGYF1/PERQ1 and GIGYF2/TNRC15., tissue specificity: Highly expressed in skeletal muscle.,