

Product name:	HNF-4 α (Acetyl Lys106) Rabbit Polyclonal Antibody
Cat number:	ABN06219
Conjugate:	Unconjugated
Size:	100 μ L
Clone:	Polyclonal
Concentration:	1mg/ml
Host:	Rabbit
Isotype:	IgG
Immunogen:	Synthetic Acetyl peptide from human protein at AA range: 106
Reactivity:	Human,Rat,Mouse
Applications:	WB 1:500-1:2000,ELISA 1:10000-1:20000
Molecular Weight:	55kDa
Purification:	Affinity purification
Form:	Liquid
Buffer:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Storage:	Store at 4°C short term. Aliquot and store at -20°C for 12 months. Avoid freeze/thaw cycles.

Background:

The protein encoded by this gene is a nuclear transcription factor which binds DNA as a homodimer. The encoded protein controls the expression of several genes, including hepatocyte nuclear factor 1 alpha, a transcription factor which regulates the expression of several hepatic genes. This gene may play a role in development of the liver, kidney, and intestines. Mutations in this gene have been associated with monogenic autosomal dominant non-insulin-dependent diabetes mellitus type I. Alternative splicing of this gene results in multiple transcript variants encoding several different isoforms. [provided by RefSeq, Apr 2012], alternative products: Additional isoforms seem to exist, disease: Defects in HNF4A are the cause of maturity onset diabetes of the young type 1 (MODY1) [MIM:125850]; also shortened MODY-1. MODY [MIM:606391] is a form of diabetes that is characterized by an autosomal dominant mode of inheritance, onset in childhood or early adulthood (usually before 25 years of age) and a primary defect in insulin secretion. The clinical phenotype of MODY1 is characterized by severe insulin secretory defects, and by major hyperglycemia associated with microvascular complications., function: Transcriptionally controlled transcription factor. Binds to DNA sites required for the transcription of alpha 1-antitrypsin, apolipoprotein CIII, transthyretin genes and HNF1-alpha. May be essential for development of the liver, kidney and intestine., miscellaneous: Binds fatty acids., online information: Hepatocyte nuclear factors entry, PTM: Phosphorylated on tyrosine residue(s); phosphorylation is important for its DNA-binding activity. Phosphorylation may directly or indirectly play a regulatory role in the subnuclear distribution., similarity: Belongs to the nuclear hormone receptor family., similarity: Belongs to the nuclear hormone receptor family. NR2 subfamily., similarity: Contains 1 nuclear receptor DNA-binding domain., subunit: Homodimerization is required for HNF4-alpha to bind to its recognition site.,